

# Random walks and orthogonal polynomials: some challenges

F. Alberto Grünbaum

**Abstract.** The study of several naturally arising "nearest neighbours" random walks benefits from the study of the associated orthogonal polynomials and their orthogonality measure. I consider extensions of this approach to a larger class of random walks. This raises a number of open problems.

To Henry, teacher and friend, with gratitude and admiration.

## 1. Introduction

Consider a birth and death process, i.e., a discrete time Markov chain on the nonnegative integers, with a one step transition probability matrix  $\mathbb{P}$ . There is then a time-honored way of writing down the  $n$ -step transition probability matrix  $\mathbb{P}^n$  in terms of the orthogonal polynomials associated to  $\mathbb{P}$  and the spectral measure. This goes back to [KMcG] and, as they observe, this is nothing but an application of the spectral theorem. One can find some precursors of these powerful ideas, see for instance [H, LR]. In as much as this is such a deep and general result, it holds in many setups, such as a nearest neighbours random walk on the  $N$ th-roots of unity. In general this representation of  $\mathbb{P}^n$  allows one to relate properties of the Markov chain, such as recurrence or other limiting behaviour, to properties of the orthogonality measure.

In the few cases when one can get one's hands on the orthogonality measure and the polynomials themselves this gives fairly explicit answers to various questions.

---

2000 Mathematics Subject Classification. 33C45, 22E45.

Key words and phrases. Matrix valued orthogonal polynomials, Karlin-McGregor representation, Jacobi polynomials.

The author was supported in part by NSF Grant # DMS 0603901.

The two main drawbacks to the applicability of this representation (to be recalled below) are:

- a) typically one cannot get either the polynomials or the measure explicitly.
- b) the method is restricted to “nearest neighbour” transition probability chains that give rise to tridiagonal matrices and thus to orthogonal polynomials.

The challenge that we pose in this paper is very simple: try to extend the class of Markov processes whose study can benefit from a similar association.

There is an important collection of papers that study in detail the cases where the entries in  $\mathbb{P}$  depends linearly, quadratically or even rationally on the index  $n$ . We make no attempt to review these results, but we just mention that the linear case involves (associated) Laguerre and Meixner polynomials, and the other cases involve associated dual Hahn polynomials. For a very small sample of important sources dealing with this connection see [Ch, vD, ILMV].

The plan for this paper is as follows:

- 1) we first review briefly the approach of S. Karlin and J. McGregor. This is done in section 2.
- 2) we consider a few examples of physically important Markov chains that happen to feature rather well known families of orthogonal polynomials. This is done in sections 3,4,5.
- 3) we propose a way of extending this representation to the case of certain Markov chains where the one-step transition probability matrix is not necessarily tridiagonal. For concreteness we restrict ourselves to the case of pentadiagonal matrices or more generally block tridiagonal matrices. This is illustrated with some examples. This material takes up sections 6,7,8,9 and 10.
- 4) A number of open problems are mentioned along the way. A few more are listed in section 11.

Both parts 1) and 2) above are well known while the proposal in 3) appears to be new.

## 2. The Karlin-McGregor representation

If we have

$$\mathbb{P}_{i,j} = Pr\{X(n+1) = j | X(n) = i\}$$

for the 1-step transition probability of our Markov chain, and we put  $p_i = \mathbb{P}_{i,i+1}$ ,  $q_{i+1} = \mathbb{P}_{i+1,i}$ , and  $r_i = \mathbb{P}_{i,i}$  we get for the matrix  $\mathbb{P}$ , in the case of a birth and death process, the expression

$$\mathbb{P} = \begin{pmatrix} r_0 & p_0 & 0 & 0 & \\ q_1 & r_1 & p_1 & 0 & \\ 0 & q_2 & r_2 & p_2 & \\ & & \ddots & \ddots & \ddots \end{pmatrix}$$

We will assume that  $p_j > 0$ ,  $q_{j+1} > 0$  and  $r_j \geq 0$  for  $j \geq 0$ . We also assume  $p_j + r_j + q_j = 1$  for  $j \geq 1$  and by putting  $p_0 + r_0 \leq 1$  we allow for the state  $j = 0$  to be an absorbing state (with probability  $1 - p_0 - r_0$ ). Some of these conditions can be relaxed.

If one introduces the polynomials  $Q_j(x)$  by the conditions  $Q_{-1}(0) = 0$ ,  $Q_0(x) = 1$  and using the notation

$$Q(x) = \begin{pmatrix} Q_0(x) \\ Q_1(x) \\ \vdots \end{pmatrix}$$

we insist on the recursion relation

$$\mathbb{P}Q(x) = xQ(x)$$

one proves the existence of a unique measure  $\psi(dx)$  supported in  $[-1, 1]$  such that

$$\pi_j \int_{-1}^1 Q_i(x) Q_j(x) \psi(dx) = \delta_{ij}$$

and one gets the Karlin-McGregor representation formula

$$(\mathbb{P}^n)_{ij} = \pi_j \int_{-1}^1 x^n Q_i(x) Q_j(x) \psi(dx).$$

Many general results can be obtained from the representation formula given above. Some of these will be given for certain examples in the next three sections.

Here we just remark that the existence of

$$\lim_{n \rightarrow \infty} (\mathbb{P}^n)_{ij}$$

is equivalent to  $\psi(dx)$  having no mass at  $x = -1$ . If this is the case this limit is positive exactly when  $\psi(dx)$  has some mass at  $x = 1$ .

If one notices that  $Q_n(x)$  is nothing but the determinant of the  $(n+1) \times (n+1)$  upper-left corner of the matrix  $xI - \mathbb{P}$ , divided by the factor

$$p_0 p_1 \cdots p_{n-1},$$

and one defines the polynomials  $q_n(x)$  by solving the same three-term recursion relation satisfied by the polynomials  $Q_n(x)$ , but with the indices shifted by one, and the initial conditions  $q_0(x) = 1$ ,  $q_1(x) = (x - r_1)/p_1$  then it is clear that the  $(0, 0)$  entry of the matrix

$$(xI - \mathbb{P})^{-1}$$

should be given, except by the constant  $p_0$ , by the limit of the ratio

$$q_{n-1}(x)/Q_n(x).$$

On the other hand the same spectral theorem alluded to above establishes an intimate relation between

$$\lim_{n \rightarrow \infty} q_{n-1}(x)/Q_n(x)$$

and

$$\int_{-1}^1 \frac{d\psi(\lambda)}{x - \lambda}.$$

We will see in some of the examples a probabilistic interpretation for the right hand side of the expression above in terms of generating functions.

The same connection with orthogonal polynomials holds in the case of a birth and death process with continuous time, and this has been extensively described in the literature. The discrete time situation discussed above is enough to illustrate the power of this method.

### 3. The Ehrenfest urn model

Consider the case of a Markov chain in the finite state space  $0, 1, 2, \dots, 2N$  where the matrix  $\mathbb{P}$  given by

$$\begin{pmatrix} 0 & 1 & & & & \\ \frac{1}{2N} & 0 & \frac{2N-1}{2N} & & & \\ & \frac{2}{2N} & 0 & \frac{2N-2}{2N} & & \\ & & \ddots & 0 & \ddots & \\ & & & \ddots & \ddots & \ddots \\ & & & & \ddots & 0 & \frac{1}{2N} \\ & & & & & \frac{2N}{2N} & 0 \end{pmatrix}$$

This situation arises in a model introduced by P. and T. Ehrenfest, see [E], in an effort to illustrate the issue that irreversibility and recurrence can coexist. The background here is, of course, the famous  $H$ -theorem of L. Boltzmann.

For a more detailed discussion of the model see [F, K]. This model has also been considered in dealing with a quantum mechanical version of a discrete harmonic oscillator by Schroedinger himself, see [SchK].

In this case the corresponding orthogonal polynomials (on a finite set) can be given explicitly. Consider the so called Krawtchouk polynomials, given by means of the (truncated) Gauss series

$${}_2\tilde{F}_1\left(\begin{matrix} a, b \\ c \end{matrix}; z\right) = \sum_0^{2N} \frac{(a)_n(b)_n}{n!(c)_n} z^n$$

with

$$(a)_n \equiv a(a+1)\dots(a+n-1), \quad (a)_0 = 1$$

The polynomials are given by

$$\begin{aligned} K_i(x) &= {}_2\tilde{F}_1\left(\begin{matrix} -i, -x \\ -2N \end{matrix}; 2\right) \\ x &= 0, 1, \dots, 2N; \quad i = 0, 1, \dots, 2N \end{aligned}$$

Observe that

$$K_0(x) \equiv 1, K_i(2N) = (-1)^i.$$

The orthogonality measure is read off from

$$\sum_{x=0}^{2N} K_i(x)K_j(x) \frac{\binom{2N}{x}}{2^{2N}} = \frac{(-1)^i i!}{(-2N)_i} \delta_{ij} \equiv \pi_i^{-1} \delta_{ij} \quad 0 \leq i, j \leq 2N.$$

These polynomials satisfy the second order difference equation

$$\frac{1}{2}(2N-i)K_{i+1}(x) - \frac{1}{2}2NK_i(x) + \frac{i}{2}K_{i-1}(x) = -xK_i(x)$$

and this has the consequence that

$$\begin{pmatrix} 0 & 1 & & & \\ \frac{1}{2N} & 0 & \frac{2N-1}{2N} & & \\ & \frac{2}{2N} & 0 & \frac{2N-2}{2N} & \\ & & \ddots & 0 & \ddots \\ & & & \ddots & \ddots & \ddots \\ & & & & \ddots & 0 & \frac{1}{2N} \\ & & & & & \frac{2N}{2N} & 0 \end{pmatrix} \begin{pmatrix} K_0(x) \\ K_1(x) \\ \\ \\ \\ K_{2N}(x) \end{pmatrix} = \left(1 - \frac{x}{N}\right) \begin{pmatrix} K_0(x) \\ K_1(x) \\ \vdots \\ K_{2N}(x) \end{pmatrix}$$

any time that  $x$  is one of the values  $x = 0, 1, \dots, 2N$ . This means that the eigenvalues of the matrix  $\mathbb{P}$  above are given by the values of  $1 - \frac{x}{N}$  at these values of  $x$ , i.e.,

$$1, 1 - \frac{1}{N}, \dots, -1$$

and that the corresponding eigenvectors are the values of  $[K_0(x), K_1(x), \dots, K_{2N}(x)]^T$  at these values of  $x$ .

Since the matrix  $\mathbb{P}$  above is the one step transition probability matrix for our urn model we conclude that

$$(\mathbb{P}^n)_{ij} = \pi_j \sum_{x=0}^{2N} \left(1 - \frac{x}{N}\right)^n K_i(x) K_j(x) \frac{\binom{2N}{x}}{2^{2N}}.$$

We can use these expressions to rederive some results given in [K].

We have

$$(\mathbb{P}^n)_{00} = \sum_{x=0}^{2N} \left(1 - \frac{x}{N}\right)^n \frac{\binom{2N}{x}}{2^{2N}}$$

and the “generating function” for these probabilities, defined by

$$U(z) \equiv \sum_{n=0}^{\infty} z^n (\mathbb{P}^n)_{00}$$

becomes

$$U(z) = \sum_{x=0}^{2N} \frac{N}{N(1-z) + xz} \frac{\binom{2N}{x}}{2^{2N}}.$$

In particular  $U(1) = \infty$  and then the familiar “renewal equation”, see [F] given by

$$U(z) = F(z)U(z) + 1$$

where  $F(z)$  is the generating function for the probabilities  $f_n$  of returning from state 0 to state 0 for the first time in  $n$  steps

$$F(z) = \sum_{n=0}^{\infty} z^n f_n$$

gives

$$F(z) = 1 - \frac{1}{U(z)}$$

Therefore we have  $F(1) = 1$ , indicating that one returns to state 0 with probability one in finite time.

These results allow us to compute the expected time to return to state 0. This expected value is given by  $F'(1)$ , and we have

$$F'(z) = \frac{U'(z)}{U^2(z)}.$$

Since

$$U'(z) = \sum_{x=0}^{2N} \frac{N(N-x)}{(N(1-z) + xz)^2} \frac{\binom{2N}{x}}{2^{2N}}$$

we get  $F'(1) = 2^{2N}$ . The same method shows that any state  $i = 0, \dots, 2N$  is also recurrent and that the expected time to return to it is given by

$$\frac{2^{2N}}{\binom{2N}{i}}.$$

The moral of this story is clear: if  $i = 0$  or  $2N$ , or if  $i$  is close to these values, i.e., we start from a state where most balls are in one urn it will take on average a huge amount of time to get back to this state.

If on the other hand  $i = N$ , i.e., we are starting from a very balanced state, then we will (on average) return to this state fairly soon. Thus we see how the issues of irreversibility and recurrence are rather subtle.

In a very precise sense these polynomials are discrete analogs of those of Hermite in the case of the real line. For interesting material regarding this section the reader should consult [A].

#### 4. A Chebyshev type example

The example below illustrates nicely how certain recurrence properties of the process are related to the presence of point masses in the orthogonality measure. This is seen by comparing the two integrals at the end of the section.

Consider the matrix

$$\mathbb{P} = \begin{pmatrix} 0 & 1 & 0 & & \\ q & 0 & p & & \\ 0 & q & 0 & p & \\ & & \ddots & \ddots & \ddots \end{pmatrix}$$

with  $0 \leq p \leq 1$  and  $q = 1 - p$ . We look for polynomials  $Q_j(x)$  such that

$$Q_{-1}(x) = 0, \quad Q_0(x) = 1$$

and if  $Q(x)$  denotes the vector

$$Q(x) \equiv \begin{pmatrix} Q_0(x) \\ Q_1(x) \\ Q_2(x) \\ \vdots \end{pmatrix}$$

we ask that we should have

$$\mathbb{P}Q(x) = xQ(x).$$

The matrix  $\mathbb{P}$  can be conjugated into a symmetric one and in this fashion one can find the explicit expression for these polynomials.

We have

$$Q_j(x) = \left(\frac{q}{p}\right)^{j/2} \left[ (2-2p)T_j\left(\frac{x}{2\sqrt{pq}}\right) + (2p-1)U_j\left(\frac{x}{2\sqrt{pq}}\right) \right]$$

where  $T_j$  and  $U_j$  are the Chebyshev polynomials of the first and second kind.

If  $p \geq 1/2$  we have

$$\left(\frac{p}{1-p}\right)^n \int_{-\sqrt{4pq}}^{\sqrt{4pq}} Q_n(x)Q_m(x) \frac{\sqrt{4pq-x^2}}{1-x^2} dx = \delta_{nm} \begin{cases} 2(1-p)\pi, & n=0 \\ 2p(1-p)\pi, & n \geq 1 \end{cases}$$

while if  $p \leq 1/2$  we get a new phenomenon, namely the presence of point masses in the spectral measure

$$\begin{aligned} & \left(\frac{p}{1-p}\right)^n \left[ \int_{-\sqrt{4pq}}^{\sqrt{4pq}} Q_n(x)Q_m(x) \frac{\sqrt{4pq-x^2}}{1-x^2} dx \right. \\ & \quad \left. + (2-4p)\pi[Q_n(1)Q_m(1) + Q_n(-1)Q_m(-1)] \right] \\ & = \delta_{nm} \begin{cases} 2(1-p)\pi, & n=0 \\ 2p(1-p)\pi, & n \geq 1 \end{cases} \end{aligned}$$

From a probabilistic point of view these results are very natural.

## 5. The Hahn polynomials, Laplace and Bernoulli

As has been pointed out before, a limitation of this method is given by the sad fact that given the matrix  $\mathbb{P}$  very seldom can one write down the corresponding polynomials and their orthogonality measure. In general there is no reason why "physically interesting" Markov chains



will give rise to situations where these mathematical objects can be found explicitly.

The example below shows that one can get lucky: there is a very old model of the exchange of heat between two bodies going back to Laplace and Bernoulli, see [F], page 378. It turns out that in this case the corresponding orthogonal polynomials can be determined explicitly.

The Bernoulli–Laplace model for the exchange of heat between two bodies consists of two urns, labeled 1 and 2. Initially there are  $W$  white balls in urn 1 and  $B$  black balls in urn 2. The transition mechanism is as follows: a ball is picked from each urn and these two balls are switched. It is natural to expect that eventually both urns will have a nice mixture of white and black balls.

The state of the system at any time is described by  $w$ , defined to be the number of white balls in urn 1. It is clear that we have, for  $w = 0, 1, \dots, W$

$$\begin{aligned}\mathbb{P}_{w,w+1} &= \frac{W-w}{W} \frac{W-w}{B} \\ \mathbb{P}_{w,w-1} &= \frac{w}{W} \frac{B-W+w}{B} \\ \mathbb{P}_{w,w} &= \frac{w}{W} \frac{W-w}{B} + \frac{W-w}{W} \frac{B-W+w}{B}.\end{aligned}$$

Notice that

$$\mathbb{P}_{w,w-1} + \mathbb{P}_{w,w} + \mathbb{P}_{w,w+1} = 1.$$

Introduce now the dual Hahn polynomials by means of

$$\begin{aligned}R_n(\lambda(x)) &= {}_3\tilde{F}_2 \left( \begin{matrix} -n, -x, x-W-B-1 \\ -W, -W \end{matrix} \middle| 1 \right) \\ n &= 0, \dots, W; \quad x = 0, \dots, W.\end{aligned}$$

These polynomials depend in general on one more parameter.

Notice that these are polynomials of degree  $n$  in

$$\lambda(x) \equiv x(x-W-B-1).$$

One has

$$\mathbb{P}_{w,w-1}R_{w-1} + \mathbb{P}_{w,w}R_w + \mathbb{P}_{w,w+1}R_{w+1} = \left(1 - \frac{x(B+W-x+1)}{BW}\right) R_w.$$

This means that for each value of  $x = 0, \dots, W$  the vector

$$\begin{pmatrix} R_0(\lambda(x)) \\ R_1(\lambda(x)) \\ \vdots \\ R_W(\lambda(x)) \end{pmatrix}$$

is an eigenvector of the matrix  $\mathbb{P}$  with eigenvalue  $1 - \frac{x(B+W-x+1)}{BW}$ . The relevant orthogonality relation is given by

$$\pi_j \sum_{x=0}^W R_i(\lambda(x)) R_j(\lambda(x)) \mu(x) = \delta_{ij}$$

with

$$\begin{aligned} \mu(x) &= \frac{w!(-w)_x(-w)_x(2x-W-B-1)}{(-1)^{x+1}x!(-B)_x(x-W-B-1)_{w+1}} \\ \pi_j &= \frac{(-w)_j}{j!} \frac{(-B)_{w-j}}{(w-j)!}. \end{aligned}$$

The Karlin–McGregor representation gives

$$(\mathbb{P}^n)_{ij} = \pi_j \sum_{x=0}^W R_i(\lambda(x)) R_j(\lambda(x)) e^n(x) \mu(x)$$

with  $e(x) = 1 - \frac{x(B+W-x+1)}{BW}$ .

These results can be used, once again, to get some quantitative results on this process.

Interestingly enough, these polynomials were considered in great detail by S. Karlin and J. McGregor, [KMcG2] and used by these authors in the context of a model in genetics describing fluctuations of gene frequency under the influence of mutation and selection. The reader will find useful remarks in [DS].

## 6. The classical orthogonal polynomials and the bispectral problem

The examples discussed above illustrate the following point: quite often the orthogonal polynomials that are associated with important Markov chains belong to the small class of polynomials usually referred to as classical. By this one means that they satisfy not only three term recursion relations but that they are also the common eigenfunctions of some fixed (usually second order) differential operator. The search for polynomials of this kind goes back at least to [B]. In fact this issue is even older, and was considered in a paper in 1884 by Routh. See [I, R] for a more complete discussion.

In the context where both variables are continuous, this problem has been raised in [DG]. For a view of some related subjects see [HK]. The reader will find useful material in [AW, AAR, I].

## 7. Matrix valued orthogonal polynomials

Here we recall a notion due to M. G. Krein, see [K1, K2]. Given a self adjoint positive definite matrix valued smooth weight function  $W(x)$  with finite moments, we can consider the skew symmetric bilinear form defined for any pair of matrix valued polynomial functions  $P(x)$  and  $Q(x)$  by the numerical matrix

$$(P, Q) = (P, Q)_W = \int_{\mathbb{R}} P(x)W(x)Q^*(x)dx,$$

where  $Q^*(x)$  denotes the conjugate transpose of  $Q(x)$ . By the usual construction this leads to the existence of a sequence of matrix valued orthogonal polynomials with non singular leading coefficient.

Given an orthogonal sequence  $\{P_n(x)\}_{n \geq 0}$  one gets by the usual argument a three term recursion relation

$$(1) \quad xP_n(x) = A_nP_{n-1}(x) + B_nP_n(x) + C_nP_{n+1}(x),$$

where  $A_n$ ,  $B_n$  and  $C_n$  are matrices and the last one is nonsingular.

We now turn our attention to an important class of orthogonal polynomials which we will call classical matrix valued orthogonal polynomials. Very much as in [D, GPT2, GPT3] we say that the weight function is classical if there exists a second order ordinary differential operator  $D$  with matrix valued polynomial coefficients  $A_j(x)$  of degree less or equal to  $j$  of the form

$$(2) \quad D = A_2(x)\frac{d^2}{dx^2} + A_1(x)\frac{d}{dx} + A_0(x),$$

such that for an orthogonal sequence  $\{P_n\}$ , we have

$$(3) \quad DP_n^* = P_n^* \Lambda_n,$$


where  $\Lambda_n$  is a real valued matrix. This form of the eigenvalue equation (3) appears naturally in [GPT1] and differs only superficially with the form used in [D], where one uses right handed differential operators.

During the last few years there has been quite a bit of activity centered around an effort to produce families of matrix valued orthogonal polynomials that would satisfy differential equations as those above. One of the examples that have resulted from this search, see [G], will be particularly useful later on.


## 8. Pentadiagonal matrices and matrix valued orthogonal polynomials

Given a pentadiagonal scalar matrix it is often useful to think of it either in its original unblocked form or as being made, let us say, of  $2 \times 2$  blocks. These two ways of seeing a matrix, and the fact that matrix operations like multiplication can be performed "by blocks" has proved very important in the development of fast algorithms.

In the case of a birth and death process it is useful to think of a graph like




Suppose that we are dealing with a more complicated Markov chain in the same probability space, where the elementary transitions can go beyond "nearest neighbours". In such a case the graph may look as follows



The matrix  $\mathbb{P}$  going with the graph above is now pentadiagonal. By thinking of it in the manner mentioned above we get a block tridiagonal matrix. As an extra bonus, its off-diagonal blocks are triangular.

The graph below



clearly corresponds to a general block tridiagonal matrix, with blocks of size  $2 \times 2$ .

If  $\mathbb{P}_{i,j}$  denotes the  $i, j$  block of  $\mathbb{P}$  we can generate a sequence of  $2 \times 2$  matrix valued polynomials  $Q_i(t)$  by imposing the three term recursion of section 8. By using the notation of section 2, we would have

$$\mathbb{P}Q(x) = xQ(x)$$

where the entries of the column vector  $Q(x)$  are now  $2 \times 2$  matrices.

Proceeding as in the scalar case, this relation can be iterated to give

$$\mathbb{P}^n Q(x) = x^n Q(x)$$

and if we assume the existence of a weight matrix  $W(x)$  as in section 7, with the property

$$(Q_j, Q_j)\delta_{i,j} = \int_{\mathbb{R}} Q_i(x)W(x)Q_j^*(x)dx,$$

it is then clear that one can get an expression for the  $(i, j)$  entry of the block matrix  $\mathbb{P}^n$  that would look exactly as in the scalar case, namely

$$(\mathbb{P}^n)_{ij}(Q_j, Q_j) = \int x^n Q_i(x)W(x)Q_j^*(x)dx.$$

Just as in the scalar case, this expression becomes useful when we can get our hands on the matrix valued polynomials  $Q_i(x)$  and the orthogonality measure  $W(x)$ . Notice that we have not discussed

conditions on the matrix  $\mathbb{P}$  to give rise to such a measure. For this issue the reader can consult [DP, D1] and the references in these papers.

The spectral theory of a scalar double-infinite tridiagonal matrix leads naturally to a  $2 \times 2$  semi-infinite matrix. This has been looked at in terms of random walks in [P]. In [ILMV] this work is elaborated further to get a formula that could be massaged to look like the right hand side of the one above.

### 9. An explicit example

Consider the matrix valued polynomials given by the 3-term recursion relation

$$A_n \Phi_{n-1}(x) + B_n \Phi_n(x) + C_n \Phi_{n+1}(x) = t \Phi_n(x), n \geq 0$$

$$\Phi_{-1}(x) = 0, \quad \Phi_0(t) = I$$

where the entries in  $A_n, B_n, C_n$  are given below.

Notice that the matrices  $A_n$  and  $C_n$  are upper and lower triangular respectively.

$$\begin{aligned}
A_n^{11} &:= \frac{n(\alpha + n)(\beta + 2\alpha + 2n + 3)}{(\beta + \alpha + 2n + 1)(\beta + \alpha + 2n + 2)(\beta + 2\alpha + 2n + 1)} \\
A_n^{12} &:= \frac{2n(\beta + 1)}{(\beta + 2n + 1)(\beta + \alpha + 2n + 2)(\beta + 2\alpha + 2n + 1)} \\
A_n^{21} &:= 0 \\
A_n^{22} &:= \frac{n(\alpha + n + 1)(\beta + 2n + 3)}{(\beta + 2n + 1)(\beta + \alpha + 2n + 2)(\beta + \alpha + 2n + 3)} \\
B_n^{11} &:= 1 + \frac{n(\beta + n + 1)(\beta + 2n - 1)}{(\beta + 2n + 1)(\beta + \alpha + 2n + 1)} \\
&\quad - \frac{(n + 1)(\beta + n + 2)(\beta + 2n + 1)}{(\beta + 2n + 3)(\beta + \alpha + 2n + 3)} \\
&\quad - \frac{2(\beta + 1)^2}{(\beta + 2n + 1)(\beta + 2n + 3)(\beta + 2\alpha + 2n + 3)} \\
B_n^{12} &:= \frac{2(\beta + 1)(\alpha + \beta + n + 2)}{(\beta + 2n + 3)(\beta + \alpha + 2n + 2)(\beta + 2\alpha + 2n + 3)} \\
B_n^{21} &:= \frac{2(\alpha + n + 1)(\beta + 1)}{(\beta + 2n + 1)(\beta + \alpha + 2n + 3)(\beta + 2\alpha + 2n + 3)} \\
B_n^{22} &:= 1 + \frac{n(\beta + n + 1)(\beta + 2n + 3)}{(\beta + 2n + 1)(\beta + \alpha + 2n + 2)} \\
&\quad - \frac{(n + 1)(\beta + n + 2)(\beta + 2n + 5)}{(\beta + 2n + 3)(\beta + \alpha + 2n + 4)} \\
&\quad + \frac{2(\beta + 1)^2}{(\beta + 2n + 1)(\beta + 2n + 3)(\beta + 2\alpha + 2n + 3)}
\end{aligned}$$

and finally for the entries of  $C_n$  we have

$$\begin{aligned}
C_n^{11} &:= \frac{(\beta + n + 2)(\beta + 2n + 1)(\beta + \alpha + n + 2)}{(\beta + 2n + 3)(\beta + \alpha + 2n + 2)(\beta + \alpha + 2n + 3)} \\
C_n^{12} &:= 0 \\
C_n^{21} &:= \frac{2(\beta + 1)(\beta + n + 2)}{(\beta + 2n + 3)(\beta + \alpha + 2n + 3)(\beta + 2\alpha + 2n + 5)} \\
C_n^{22} &:= \frac{(\beta + n + 2)(\beta + \alpha + n + 3)(\beta + 2\alpha + 2n + 3)}{(\beta + \alpha + 2n + 3)(\beta + \alpha + 2n + 4)(\beta + 2\alpha + 2n + 5)}.
\end{aligned}$$

If the matrix  $\Psi_0(x)$  is given by

$$\Psi_0(x) = \begin{pmatrix} 1 & 1 \\ 1 & \frac{(\beta+2\alpha+3)x}{\beta+1} - \frac{2(\alpha+1)}{\beta+1} \end{pmatrix}$$

one can see that the polynomials  $\Phi_n(x)$  satisfy the orthogonality relation

$$\int_0^1 \Phi_i(x) W(x) \Phi_j^*(x) dx = 0 \text{ if } i \neq j$$

where

$$W(x) = \Psi_0(x) \begin{pmatrix} (1-x)^\beta x^{\alpha+1} & 0 \\ 0 & (1-x)^\beta x^\alpha \end{pmatrix} \Psi_0^*(x).$$

The polynomials  $\Phi_n(x)$  are “classical” in the sense that they are eigenfunctions of a fixed second order differential operator. More precisely, we have

$$\mathcal{F} \Phi_n^* = \Phi_n^* \Lambda_n$$

where

$$\Lambda_n = \begin{pmatrix} -n^2 - (\alpha + \beta + 2)n + \alpha + 1 + \frac{\beta+1}{2} & 0 \\ 0 & -n^2 - (\alpha + \beta + 3)n \end{pmatrix}$$

and  $\mathcal{F}$  is given by

$$\begin{aligned} \mathcal{F} &= x(1-x) \left( \frac{d}{dx} \right)^2 \\ &+ \begin{pmatrix} \frac{(\alpha+1)(\beta+2\alpha+5)}{\beta+2\alpha+3} - (\alpha + \beta + 3)x & \frac{\frac{2\alpha+2}{2\alpha+\beta+3} + x}{\frac{\beta+1}{\beta+2\alpha+3}} \\ \frac{\beta+1}{\beta+2\alpha+3} & \frac{(\alpha+2)\beta+2\alpha^2+5\alpha+4}{\beta+2\alpha+3} - (\alpha + \beta + 4)x \end{pmatrix} \frac{d}{dx} \\ &+ \begin{pmatrix} \alpha + 1 + \frac{\beta+1}{2} & 0 \\ 0 & 0 \end{pmatrix} I. \end{aligned}$$

As mentioned earlier this is the reason why this example has surfaced recently, see [G]. An explicit expression for the polynomials themselves is given in [T2], corollary 3.


Now we make two observations

- 1) the entries of the corresponding pentadigonal matrix are all non-negative.
- 2) the sum of the entries on any given row are all equal to 1.

This allows for an immediate probabilistic interpretation of the pentadiagonal matrix as the one step transition probability matrix for a



Markov chain whose state space could be visualized by the graph given below.



I find it rather remarkable that this example which was produced for an entirely different purpose should have this extra property. Finding an appropriate combinatorial mechanism, maybe in terms of urns, that goes along with this example remains an interesting challenge.

Two final observations dealing with these state spaces that can be analyzed using matrix valued orthogonal polynomials. If we were using matrix valued polynomials of size  $N$  we would have as state space a semiinfinite network consisting of  $N$  (instead of two) parallel collection of non-negative integers with connections going from each of the  $N$  states on each vertical rung to every one in the same rung and the two neighbouring ones. The examples in [GPT1] give instances of this situation with a rather local connection pattern.

In the case of  $N = 2$  one could be tempted to paraphrase a well known paper and say that "it has not escaped our notice that" some of these models could be used to study transport phenomena along a DNA segment.

#### 10. Another example

Here we consider a different example of matrix valued orthogonal polynomials whose block tridiagonal matrix can be seen as a scalar pentadiagonal matrix with non-negative elements. In this case the sum of the elements in the rows of this scalar matrix is not identically one, but this poses no problem in terms of a Karlin–McGregor type representation formula for the entries of the powers  $\mathbb{P}^n$ .

This example has the important property that the orthogonality weight matrix  $W(x)$ , as well as the polynomials themselves are explicitly known. This is again, a classical situation, see [CG].

Consider the block tridiagonal matrix

$$\begin{pmatrix} B_0 & I & & & \\ A_1 & B_1 & I & & \\ 0 & A_2 & B_2 & I & \\ & & \ddots & \ddots & \ddots \end{pmatrix}$$

with  $2 \times 2$  blocks given as follows

$$\begin{aligned} B_0 &= \frac{1}{2}I, \quad B_n = 0 \text{ if } n \geq 1 \\ A_n &= \frac{1}{4}I \text{ if } n \geq 1. \end{aligned}$$

In this case one can compute explicitly the matrix valued polynomials  $P_n$  given by

$$A_n P_{n-1}(x) + B_n P_n(x) + P_{n+1}(x) = x P_n(x), \quad P_{-1}(x) = 0, \quad P_0(x) = I.$$

One gets

$$P_n(x) = \frac{1}{2^n} \begin{pmatrix} U_n(x) & -U_{n-1}(x) \\ -U_{n-1}(x) & U_n(x) \end{pmatrix}$$

where  $U_n(x)$  are the Chebyshev polynomials of the second kind.

The orthogonality measure is read off from the identity

$$\frac{4^i}{\pi} \int_{-1}^1 P_i(x) \frac{1}{\sqrt{1-x^2}} \begin{pmatrix} 1 & x \\ x & 1 \end{pmatrix} P_j(x) dx = \delta_{ij} I.$$

We get, for  $n = 0, 1, 2, \dots$

$$\frac{4^i}{\pi} \int_{-1}^1 x^n P_i(x) \frac{1}{\sqrt{1-x^2}} \begin{pmatrix} 1 & x \\ x & 1 \end{pmatrix} P_j(x) dx = (\mathbb{P}^n)_{ij}$$

where, as above,


$$(\mathbb{P}^n)_{ij}$$

stands for the  $i, j$  block of the matrix  $\mathbb{P}^n$ .

In this way one can compute the entries of the powers  $\mathbb{P}^n$  with  $\mathbb{P}^n$  thought of as a pentadiagonal matrix, namely

$$\mathbb{P} = \begin{pmatrix} 0 & \frac{1}{2} & 1 & 0 & 0 & 0 & \\ \frac{1}{2} & 0 & 0 & 1 & 0 & 0 & \ddots \\ \frac{1}{4} & 0 & 0 & 0 & 1 & 0 & \ddots \\ 0 & \frac{1}{4} & 0 & 0 & 0 & 1 & \ddots \\ & 0 & \frac{1}{4} & 0 & 0 & 0 & \ddots \\ & & 0 & \frac{1}{4} & 0 & 0 & \ddots \\ & & & \ddots & \ddots & \ddots & \ddots \end{pmatrix}.$$

This example goes along with the following graph.



### 11. A few more challenges

We have already pointed out a few challenges raised by our attempt to extend the Karlin-McGregor representation beyond its original setup. Here we list a few more open problems. The reader will undoubtedly come up with many more.

Is there a natural version of the models introduced by Bernoulli-Laplace and by P. and T. Ehrenfest whose solution features matrix valued polynomials?

Is it possible to modify the simplest Chebyshev type examples in [D1] to accomodate cases where some of the blocks in the tridiagonal matrix give either absorption or reflection boundary conditions?

One could consider the emerging class of polynomials of several variables and find here interesting instances where the state space is higher dimensional. For a systematic study of polynomials in several variables one should consult [DX] as well as the work on Macdonald polynomials of various kinds, see [M]. A look at the pioneering work of Tom Koornwinder, see for instance [Ko], is always a very good idea.

After this paper was finished I came up with two independent sources of multivariable polynomials of the type alluded to in the previous paragraph. One is the series of papers by Hoare and Rahman, see [HR1, HR2, HR3, HR4]. The other one deals with the papers [Mi] and [IX, GI].

In queueing theory one finds the notion of Quasi-Birth-and-Death processes, see [LR1, N]. Within those that are nonhomogeneous one could find examples where the general approach advocated here might be useful.

## References

- [A] Askey, R. Evaluation of Sylvester type determinants using orthogonal polynomials, *Advances in Analysis, Proceed. 4th international ISAAC Congress*, ed. H.G.W. Begehr et al, World Scientific, Singapore, (2005), 1–16.
- [AAR] Andrews, G., Askey, R., and Roy, R., *Special functions, Encyclopedia of Mathematics and its applications*, Cambridge University Press, 1999.
- [AW] Askey, R., and Wilson, J., Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, *Mem. Amer. Math. Soc.*, 54 (1985) 319.
- [B] Bochner, S., Über Sturm–Liouvillesche polynomsysteme, *Math Z.* 29 (1929), 730–736.
- [CG] Castro, M. and Grünbaum F. A., The algebra of matrix valued differential operators associated to a given family of matrix valued orthogonal polynomials: five instructive examples, *International Math. Research Notices*, 2006:7 (2006), 1–33.
- [Ch] Chihara, T., *An introduction to orthogonal polynomials*, Gordon and Breach Science Publishers, 1978.
- [vD] van Doorn, E., On associated polynomials and decay rates for birth-death processes, *J. Math. Anal. Appl.* 278, (2003), 2, 500–511.
- [DG] Duistermaat, J. J., and Grünbaum, F. A., Differential equations in the spectral parameter, *Comm. Math. Phys.* 103 (1986), 177–240.

- [D] Duran, A., Matrix inner product having a matrix symmetric second order differential operators, *Rocky Mountain Journal of Mathematics* 27, No. 2, Spring 1997.
- [D1] Duran, A. J., Ratio Asymptotics for orthogonal matrix polynomials, *J. Approximation Theory* 100 (1999), 304–344.
- [DP] Duran, A. J. and Polo, B., Gauss quadrature formulae for orthogonal matrix polynomials, *Linear Alg. Appl.* 355 (2002), 119–146.
- [DS] Diaconis, P. and Shahshahani, M., Time to reach stationarity in the Bernoulli-Laplace diffusion model, *SIAM J. Math. Anal.* 18, 208–218, 1987.
- [DX] Dunkl, C. and Xu, Y., *Orthogonal polynomials of several variables*, Cambridge University Press, 2001.
- [E] Ehrenfest, P. and Eherenfest, T. , Über zwei bekannte Einwände gegen das Boltzmannsche H-Theorem, *Physikalische Zeitschrift*, vol 8, 1907, 311–314.
- [F] Feller, W., *An introduction to Probability Theory and its Applications*, vol 1, 3rd edition, Wiley 1967.
- [GI] Geronimo, J. and Iliev, P., Two variable deformations of the Chebyshev measure arXiv.math.CA/0612489v1
- [G] Grünbaum, F. A., Matrix valued Jacobi polynomials, *Bull. Sci. Math* 127, (2003) 207–214.
- [GPT1] Grünbaum, F. A., Pacharoni, I., and Tirao, J. A., Matrix valued spherical functions associated to the complex projective plane, *J. Functional Analysis* 188 (2002) 350–441.
- [GPT2] Grünbaum, F. A., Pacharoni, I., and Tirao, J. A., Matrix valued orthogonal polynomials of the Jacobi type, *Indag. Mathem.* 14 3,4 (2003), 353–366.
- [GPT3] Grünbaum F. A., Pacharoni I. and Tirao J. A., Matrix valued orthogonal polynomials of the Jacobi type: The role of group representation theory, *Ann. Inst. Fourier, Grenoble*, 55 , 6, (2005), 2051–2068.
- [H] Harris, T. E., First passages and recurrence distributions, *Trans. American Math. Soc.* 73, (1952) 471–486.
- [HK] Harnard, J., and Kasman, A. (editors) *The bispectral problem*, CRM Proceedings and lecture notes. Univ. de Montreal, vol 14, 1998.
- [HR1] M. Hoare and M. Rahman, Distributive processes in discrete systems, *Physica*, 97 A, (1979), 1–41.
- [HR2] M. Hoare and M. Rahman, Cumulative Bernoulli trials and Krawtchouk processes, *Stochastic Processes and their applications*, 16 , (1983), 113–139
- [HR3] M. Hoare and M. Rahman, Cumulative hypergeometric processes: a statistical role for the  ${}_nF_{n-1}$  functions, *J. Math. Anal. Appl.* , 135 (2), (1988) , 615–626.
- [HR4] M. Hoare and M. Rahman, A probabilistic origin for a new class of bi-variate polynomials, to appear in *JCAM*, 2007.
- [IX] Iliev, P. and Xu, Y. , Discrete orthogonal polynomials and difference equations of several variables, arXiv.math.CA/0508039v2
- [I] Ismail, M.E.H., *Classical and Quantum Orthogonal Polynomials in One Variable*, Cambridge University Press, 2005.

- [ILMV] Ismail, M.E.H., Letessier, J., Masson, D., and Valent, G., Birth and death processes and orthogonal polynomials, *Orthogonal Polynomials*, P. Nevai (editor) Kluwer Acad. Publishers, 1990.
- [K] Kac, M., Random walk and the theory of Brownian motion, *American Math. Monthly* (1947) 54, 369–391.
- [KMcG] Karlin, S., and McGregor, J., The classification of birth and death processes, *TAMS* 86, 366–400, 1957.
- [KMcG2] Karlin, S., and McGregor, J., The Hahn polynomials, formulas and applications, *Scripta Math.* 26, 33–46.
- [Ko] Koornwinder, T. , Two variable analogues of the classical orthogonal polynomials, in *Theory and Applic. of Special Functions*, R. Askey (ed.) Academic Press, 1975, 435–495.
- [K1] Krein, M. G., Fundamental aspects of the representation theory of hermitian operators with deficiency index  $(m, m)$ , *AMS Translations, Series 2*, vol. 97, Providence, Rhode Island (1971), 75–143.
- [K2] Krein, M. G., Infinite  $J$ -matrices and a matrix moment problem, *Dokl. Akad. Nauk SSSR* 69 nr. 2 (1949) , 125–128.
- [LR1] Latouche, G., and Ramaswami, V., *Introduction to Matrix Analytic Methods in Stochastic Modeling*, ASA-SIAM Series on Statistics and Applied Probability, 1999.
- [LR] Ledermann, W., and Rueter, G. E., Spectral theory for the differential equations of simple birth and death processes, *Philos. Trans. Roy. Soc. London, Ser. A.* 246, 1954, 321–369.
- [M] Macdonald, I. , *Affine Hecke algebras and orthogonal polynomials*, Cambridge Tracts in Mathematics, vol. 157, Cambridge University Press, 2003.
- [Mi] Milch, P. , A multidimensional linear growth birth and death processes, *The Annals of Math. Statistics*, 1968 , vol. 39, 3, pp 727–754.
- [N] Neuts, M. F. , *Structured Stochastic Matrices of  $M/G/1$  Type and Their Applications*, Marcel Dekker, NY, 1989.
- [P] Pruitt, W. E. Bilateral birth and death processes, *TAMS* 107, 1963, 508–525.
- [R] Routh, E., On some properties of certain solutions of a differential equation of the second order, *Proc. London Math. Soc.*, 16 (1884), 245–261.
- [SchK] Schroedinger, E. and Kohlrausch , Das Ehrenfestche Model der H-Kurve, *Phys. Zeit.* , 8, (1907) 311–314.
- [T2] Tirao, J., The matrix valued hypergeometric equation, *Proc. Nat. Acad. Sci. U.S.A.*, 100 nr. 14 (2003), 8138–8141.

Department of Mathematics, University of California, Berkeley, CA 94720